

Application of an integrated hydrological nowcasting chain on Liguria Region

Maria Laura Poletti¹, Francesco Silvestro¹, Silvio Davolio², Flavio Pignone¹, and Nicola Rebora¹

1 CIMA Research Foundation; 2 ISAC-CNR



Introduction to nowcasting (1)

Why nowcasting is important in hydrological application?


Area of application:
Liguria Region

Catchments with
small drainage area

Short response time
(few hours!)

Forecast at short lead
time is essential!

To predict precipitation at small temporal
(few hours) and spatial (few km) scale



Mediterranean area
(Spain, France, Italy)
9-10-11 October 2018
16 victims.


Introduction to nowcasting (2)

Limit of the nowcasting models:
forecast horizon up to **few hours**



In the nowcasting models the evolution of the precipitation systems is not included: the physics representing growth and decay of the precipitation systems becomes progressively more important with increasing lead time

Connection of nowcasting and meteorological model forecasted rainfall fields

To have rainfall forecasts more accurate as possible




An integrated hydrological nowcasting chain




The nowcasting algorithm

Technique used to combine the forecasted rainfall fields in a smooth way



Observed radar rainfall field

Rainfall fields forecasted by **Numerical Weather Prediction** model corrected with Data Assimilation




The NWP model corrected with DA



Application on Liguria Region

INPUT

Hydrological model




The elements of the chain (1): the nowcasting algorithm


PhaSt

Phase Stochastic nowcasting technique

Metta et al., 2009



1. Starting point: last two rainfall maps provided by the meteorological radar



2. Gaussianization of the fields

3. Fourier transformation of the gaussianized fields

Spectral amplitudes are kept constant


Phase velocities (i.e. angular frequencies) evolve

OUTPUT
Rainfall ensemble

4. Non-linear anti-transformation of the predicted gaussian field

Generation of an ensemble of possible evolution of precipitation field

Elements of the chain (2) Numerical Weather Prediction model corrected with Data Assimilation



NWP model

MOLOCH

Malguzzi et al., 2006
Buzzi et al., 2014


non-hydrostatic; it explicitly solves deep convection

Spatial resolution ≈ 2.2 km

Integration domain north and central Italy

Initial and boundary conditions provided at 1-hour interval by the BOLAM (Limited Area Model) forecasts

Use of the **deterministic** forecast




Hourly precipitation estimates provided by Settepani radar are used as observations

DA technique

Nudging

Model specific humidity profiles at each grid point are progressively modified depending on the comparison between **observed** and forecast rainfall



Elements of the chain (3): the hydrological model

Continuous distributed hydrological model

Continuum


Silvestro et al., 2013
Silvestro et al., 2015b

It solves the hydrological processes on a longer period of time. It considers all the processes involved in the hydrologic cycle (overland and channel flow, infiltration and subsurface flows, deep flow, vegetation interception, energy balance and evapotranspiration)

The model is based on a space-filling representation of the network, directly derived from a DEM, that allows to identify flow directions on the basis of the directions of maximum slope.

OUTPUT

**Discharge
forecast**



The blending technique: a new approach

1

«Classical blending» technique: linear combination of the nowcasted rainfall fields with the NWP forecasted rainfall field corrected with DA



Linear combination obtained through **weighting** of the different fields according to the blending function

Short lead time:
more weight to
nowcasted
rainfall fields

Longer lead time:
more weight to
NWP forecasted
rainfall fields

$$\text{Rainfall field}_{\text{blended}}(T) = (\text{weight}_{\text{nowc}}(T) * \text{rain}_{\text{nowc}}(T)) + (\text{weight}_{\text{nud}}(T) * \text{rain}_{\text{nud}}(T))$$

Blending function




$$\text{Weight}_{\text{NWP}} = 1 - \text{Weight}_{\text{NOWC}}$$

The blending technique: a new approach



2

Modification of the nowcasted rainfall field with the volume trend estimated by the rainfall field forecasted with NWP corrected with DA



Results: analysis of the hydrological output

Comparison of 3 configurations:



The spread of the discharge forecast ensemble is markedly smaller when input rainfall is provided by blending (red envelope) instead of nowcasting alone (blue and orange envelopes) → smaller variance!

Results: some case studies

2 analysis

1 Punctual analysis → at basin scale

2 Distributed analysis → all over the computation domain of the hydrological model

Even if the events analysed are few the sampling points that can be analysed are thousands!

Scores used for the analysis:

Nash Sutcliffe efficiency

$$NS = 1 - \frac{\sum_{t=1}^T (Q_m(t) - Q_{obs}(t))^2}{\sum_{t=1}^T (Q_{obs} - \bar{Q}_{obs})^2}$$

Variance

$$Var(X) = E[(X - \mu)^2]$$

Reduced Continuous Rank Probability Score

$$RCRPS(F, x) = \frac{1}{\sigma^2} \int_{-\infty}^{\infty} (F(y) - \mathbb{1}(y - x))^2 dy$$

Results: punctual analysis at basin scale (1)

Case event: 9th October 2014

Bisagno creek flood
(Genova)




Nash Sutcliffe coefficient shows similar performances of the different configurations

Variance is actually smaller for the configurations with blending

RCRPS shows no clear enhancing of the performance of the chain with the use of blending

For this event, the forecast of the meteorological model, even corrected with data assimilation, is not able to improve the QPF.



Scores calculated for Bisagno at Passerella Firpo A= 97 kmq 9th October event



Results: distributed analysis (1)

9th October 2014

In this case the **RCRPS** behavior shows that the use of the information retrieved by the NWP model in the rain forecast worsen the hydrological forecast.



Particular **type of event**: stationary and persistent heavy precipitation on the same portion of territory

The event was not forecast precisely by the NWP model, but well reproduced by the nowcasting model



Results: punctual analysis at basin scale (2)

Case event: 11th November 2014

Event involving Entella basin and its tributaries (Chiavari)




Nash Sutcliffe coefficient and Variance show the same results of 9 October

RCRPS clearly highlight better performances of the configurations that are using the blending

While rainfall fields from nowcasting techniques lead to an overestimation of the discharge, the rainfall fields obtained through the blending clearly improves the discharge forecast.

Scores calculated for Graveglia at Caminata A= 42 kmq 11th November event




Results: distributed analysis (2)

11th November 2014

In this case the **RCRPS** behavior shows, as in the punctual analysis, that the configuration using the blending performs markedly better.

REDUCED CONTINUOUS RANK PROBABILITY SCORE 11th November



For this event it is also worth to note the **different behavior of the score depending on the class area of the point analyzed**

Especially for the bigger basins, due to their longer response time, the effects of a proper rainfall forecast provided with blending are beneficial for longer lead times.

Conclusions

- ❖ The use of an integrated nowcasting hydrological chain is useful in real time as a support for Civil Protection actions.
- ❖ The use of the best rainfall forecasts available at each time step can improve the hydrological forecast.
- ❖ The blending technique is useful to smoothly connect the forecasts result of nowcasting and of the NWP model but the goodness of the resulting rainfall field is really sensitive to the quality of the NWP model forecast.

TO DO:

- *Extend the analysis to other case studies*
- *Use different NWP models and DA assimilation techniques to be combined with nowcasting*
- *Explore other blending techniques*
- *Use of nowcasted rainfall fields in DA*

*Thank
you*