



Observed trends in frost and hours of cold in Majorca

Journal:	<i>International Journal of Climatology</i>
Manuscript ID:	JOC-09-0168.R1
Wiley - Manuscript type:	Short Communication
Date Submitted by the Author:	
Complete List of Authors:	Cuxart, Joan; University of the Balearic Islands, Physics Guijarro, Jose Antonio; Agencia Estatal de Meteorologia, D.T en Illes Balears
Keywords:	series of minimum temperature, Majorca, frost, hours-of-cold, cloud cover, warmer winters, agricultural damage



Only

1
2
3
4
5
6
7
8
9 Observed trends in frost and hours of cold in Majorca
10
11

12
13
14 Short title: Cold in Majorca
15
16
17

18 Authors: J. Cuxart (1*) and J.A. Guijarro (2)
19
20

21 (1) Universitat de les Illes Balears (UIB), Grup de Meteorologia
22

23 (2) Agencia Estatal de Meteorologia (AEMET), D.T. en Illes Balears
24
25
26
27

28 (*) Corresponding author address: Facultat de Ciències, Carretera de
29

30 Valldemossa km 7.5; 07122 Palma; Illes Balears. joan.cuxart@uib.cat; Ph: +34 971
31

32 171375; Fax: +34 971 173426
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract:

The observed trends on some cold indexes in the island of Majorca are analyzed using the available series of the daily minimum temperatures of the Majorca Airport (1972-2008), supplemented with the inspection of eight more climatological stations of the island covering at least thirty simultaneous years. The analysis yields a statistically significant trend to a rise of about 0.6°C per decade for the ensemble of the stations. There is also a clear shortening of the season with days with T below 0°C or 7°C, but the yearly minima or the dates of the first and last occurrence of these values do not show so well defined tendencies. A rise in nighttime cloud cover could be related to such behaviour.

Keywords: series of minimum temperature, Majorca. warmer winters, frost, hours of cold, cloud cover

Acknowledgements: The Spanish Agency of Meteorology (AEMET) has provided the time series. The Spanish Ministry of Science funds partially this work through grant CGL2006-12474-C01. M. A. Jiménez (UIB) is acknowledged for her help with figure 1.

1. Introduction

In the frame of the global warming, a number of issues are relevant locally. In this research note, the August 1972- May 2008 series of the Palma Airport -and eight more series of Majorca of 30 or more years- (Balearic Islands, Western Mediterranean) of minimum temperatures are analyzed to inspect if there is a significant trend to warmer winters and reduction of the risk of frost, mainly with agricultural intention.

A study with a similar purpose for the United States was made by Easterling (2002) with a large number of stations, that showed a significant trend to the reduction of the number of frost days at the national scale, at a rate of 0.8 days per decade for the period 1948-1999, with large regional differences. Other studies include those of Menzel *et al.* (2003), showing many stations in Germany, Austria and Estonia with significant changes in these parameters, for instance 4.9 days per decade more of the frost-free period between 1951 and 2000, or Bartolini *et al.* (2008) that show increases of more than 0.2°C per decade in the minimum temperatures for Tuscany during 1955-2004. Not all these indices are directly comparable, but they all point in the same direction. The Fourth Assessment Report of the IPCC (Trenberth *et al.*, 2007) provides estimations for land over the northern hemisphere of increases of about 0.3C per decade for the minimum temperatures (based on Vose et al, 2005) and 0.2 per°C decade for the mean temperature over ocean (based on Rayner *et al*, 2006) for the period 1979-2005, similar to the one studied here. The fact that Majorca is a piece of land in the middle of the Mediterranean sea makes interesting to look at its specific behaviour.

1
2
3
4 Majororca is the largest island of the Balearic archipelago, located in the Western
5
6 Mediterranean about 200 km offshore of the coast of the Iberian Peninsula (Fig. 1a).
7
8 The island has a characteristic size of 100 km with a large mountain range (Serra de
9
10 Tramuntana) occupying all its northwestern side (peaking at 1436 m above sea level –
11
12 ASL-). The rest of the island consists summarily in three basins, two at the southern
13
14 side (Palma at west and Campos at the east) and one at the northern side (Alcúdia), with
15
16 an elevated area in the center of the island (Randa, 556 ASL) and a lower mountain
17
18 range at the east (Serra de Llevant, peaking at 520 m ASL) -see Figure 1b. A study by
19
20 Cuxart *et al.* (2007) found that, under clear skies and with weak pressure gradients,
21
22 when strong local surface cooling may occur, the flow is organized by basins, each
23
24 having sea-breezes in the daytime and convergence of drainage flows in nighttime, with
25
26 formation of cold pools in the valleys and cold outflow to the sea. These conclusions
27
28 were reproduced by Jimenez *et al.* (2008), in a work that verified the structures in the
29
30 island using the thermal structures produced by infrared satellite imagery. The fact that
31
32 the island is relatively small and surrounded by an almost closed sea may indicate that a
33
34 warming there is regionally relevant.
35
36
37
38
39
40
41
42
43
44

45 The main agricultural cultures in the archipelago are ornamental plants, fruits (including
46
47 vines), potatoes and green vegetables. The productivity of the latter is mainly affected
48
49 by frosts whereas fruit trees have also specific requirements of hours-of-cold
50
51 (customarily defined as hours with temperature below 7°C). Any significant trend may
52
53 induce significant changes in the agriculture of the islands, thus in the economy and the
54
55 landscape, since 37% of the surface is cultivated land. To illustrate the impact of frost
56
57 episodes, all cultures of the island are listed in the insurance coverages by the spanish
58
59
60

1
2
3
4 ministry of agriculture, at the same level of continental Spain, the most sensitive sectors
5 being the early potato production for export to North Europe and the almond trees with
6 early floration in January and February. These two sectors amount respectively almost
7 one third and one tenth of the total income of the agricultural sector (Riera and Ripoll,
8 2009) whereas vegetables and flowers are usually cultivated in greenhouses . To review
9 the fundamentals of frost and how to protect the cultures against it the reader is referred
10 to Kalma *et al.* (1992) or Snyder and Melo de Abreu (2005). In this research note, the
11 series of nine stations located in diferent geomorphological areas of the island are
12 analyzed and the trends of the cold indices discussed.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27

28 2. The series

29
30
31
32
33 In Figure 1b, the locations of the nine analyzed stations are displayed. The Airport
34 Meteorological Station (WMO principal station) is located in the eastern part of the
35 Palma basin, at a distance of 9 km from downtown Palma and at 5 km of its continuous
36 urbanized area. The data are taken at the northern end of the runaway, 5 km inland, in a
37 local depression of the terrain where cold air can cumulate in clear and calm nights
38 (Cuxart *et al.*, 2007). The series starts in August 1972, complying with all the
39 requirements of a primary WMO station; it is managed and quality-controlled by the
40 Spanish Meteorological Agency (AEMET). The series of minimum temperatures from
41 1 August 1972 to 31 May 2008 has been extracted from the data base.
42
43
44
45
46
47
48
49
50
51
52
53
54
55

56 Eight other stations from the climatological database of AEMET have been used
57 provided that they had at least 30 years falling in the period 1972-2008. In the Palma
58
59
60

1
2
3
4 basin, besides the airport, there is Portopí, in the harbour of the city of Palma (1978-
5
6 2008, 3m ASL) and Sa Cabaneta, close to elevated plateau in the center of the island
7
8 (1976-2008, 152 m ASL). In the Campos basin, we could consider Lluçmajor (1974-
9
10 2008, 140 m ASL), also in an elevated part between the Palma and Campos basins, and
11
12 Ses Salines (1972-2008, 3m ASL), very near to the coast but in the center of the valley.
13
14 In the Alcúdia basin, the station of Muro is available (1974-2008, 50 m ASL) in the flat
15
16 central part of the basin, and Manacor (1974-2008, 125 m) which is more inland and to
17
18 the East. Finally, two stations are in the main Tramuntana Range, Lluç (1972-2008, 490
19
20 m ASL) and Orient (1975-2008, 455 m ASL), both located at the bottom of elevated
21
22 closed mountain valleys.
23
24
25
26
27
28
29

30 The series consist of the daily minimum temperatures, with very few missing data. The
31
32 analysis of trends for the main series has been made for complete years to avoid the
33
34 effect of starting an analysis in winter and finishing it in summer, which would provide
35
36 spurious warming. All the selected series start and finish in summer for this purpose.
37
38 To study their relative homogeneity, they were first standardized, and reference series
39
40 for each station were computed as an average of their neighbours, weighted by an
41
42 inverse distance function. Then, the differences between the original and computed
43
44 series were scrutinized in search of possible shifts in the average, by means of applying
45
46 two sample Student t-tests to two adjacent running windows of four years (48 monthly
47
48 values per sample) along all the series (in this way, four years is the maximum temporal
49
50 resolution to detect shifts either consecutive of at the beginning or the end of the series).
51
52
53
54
55
56
57
58
59
60

1
2
3
4 This lead to the detection of clear shifts in the mean in two of the series, at the
5
6 following dates: 1) March 1983 in Muro, due to a documented bad siting of the
7
8 thermometer (on a terrace); and 2) May 1984 in Manacor, for which no metadata were
9
10 found to explain this inhomogeneity. Therefore, data previous to the shifts were deleted
11
12 from the Muro and Manacor series in order to avoid misleading results. The rest of the
13
14 compared series showed minor differences, indistinguishable from natural variability,
15
16 and therefore no further corrective measures were attempted. The statistical analyses
17
18 have been done with the R statistical package (R Development Core Team, 2008),
19
20 package "Climatol" (Guijarro, 2006), version 2.0.
21
22
23
24
25
26
27
28
29
30
31
32

33 **3. Trends of minimum temperatures and cold indexes**

34
35
36
37 The trends for daily minimum series have been computed for the nine locations and the
38
39 values are given in Table 1, keeping only the restricted ranges for the two stations with
40
41 confirmed inhomogeneities. In that table the stations are ordered from warmer to colder
42
43 in respect to the average minimum temperature. The trends have been computed by the
44
45 least-squares methods and the confidence levels have been assessed with the non-
46
47 parametric Mann-Kendall test (1975), to overcome the lack of normality of the series
48
49 of the daily minimum temperatures. All the stations have positive slopes statistically
50
51 significant at least with a 95% confidence. An averaged central value for the ensemble
52
53 of the stations is 0.6°C per decade, the double of the estimation of the IPCC (Trenberth
54
55 et al) for the northern hemisphere over land. This indicates that there is a warming of
56
57
58
59
60

1
2
3
4 the nocturnal temperatures in the island at all kinds of landscape and height, since
5
6 stations are near the coast, in the valleys, in plateaux or mountain ranges. In the case of
7
8 the Airport, the complete series has a correlation coefficient of 0.11, meaning that most
9
10 of the variability is not explained by the linear regression. This is not strange, since
11
12 minima for this station can vary from -6 to 27°C and most of the variance is explained
13
14 by the annual cycle.
15
16
17

18
19
20 A number of indices have been obtained from each series, namely i) the yearly
21
22 minimum temperature value; ii) the number of days below 0°C, iii) the first day with T
23
24 below 0°C, iv) the last day below 0°C, v) the number of days below 7°C, vi) the first day
25
26 with T below 7°C, and vii) the last day below 7°C. The resulting annual series for the
27
28 Airport are plotted in Figure 2. The absolute annual minimum temperature is shown to
29
30 have a smaller trend than the annual anomaly of the same quantity. Although at a first
31
32 glance the linear adjustments seem to indicate a behaviour consistent with a general
33
34 warming, some of these results are not statistically significant and deserve further
35
36 analysis.
37
38
39
40
41
42
43

44
45 Also in Table 1 the average values and the trends for the yearly minimum temperature
46
47 are given. Average values below zero are found in the mountain range and in the bottom
48
49 of the valleys. However only half of the stations show a significant increase, making
50
51 less clear the trend to higher absolute yearly minimum temperatures in the Island.
52
53

54
55
56 Table 2 displays statistics related to frost days. Averages above 1 day per year are found
57
58 only in four stations in the mountain and in the valleys and at all these stations there is a
59
60

1
2
3
4 significant trend to diminish the number of days, with an ensemble central value of -4.7
5
6 days per decade, a trend that would indicate that frost would disappear from the valleys
7
8 in about 4 decades. However the bottom end of the 95% interval is very low for three of
9
10 the four stations (-0.2, -0.9 and -1.4 days/decade) and it suggests to be prudent in any
11
12 conclusion regarding this issue. There is not a well defined trend for the occurrence of
13
14 the first frost for these stations, although one can see that the last spring frost seems to
15
16 come earlier (about 12 days per decade).
17
18
19
20

21
22
23 Similar conclusions are reached with the same indices using days with temperatures
24
25 below 7°C (Table 3). Eight of the nine stations show a decrease in the number of days,
26
27 since Lluçmajor does not yield any significant trend. It cannot be said that the first day
28
29 with temperature below 7°C has a well defined tendency for the ensemble of stations,
30
31 but six of nine stations show statistically significant trend to earlier end of cold days.
32
33 Since some hundreds of hours below 7°C are needed by fruit trees in their winter rest,
34
35 and the average computed decrease is about 12 days below 7°C per decade, some of the
36
37 current plantations in the Balearic Islands may cease to be possible, at least with the
38
39 present-day plant varieties.
40
41
42
43
44
45
46

47 **4. Trends on cloudiness and relative insolation**

48
49
50

51
52 The fact that minimum temperatures seem to be increasing and the number of days of
53
54 frost ($T < 0^{\circ}\text{C}$) and of cold ($T < 7^{\circ}\text{C}$) decreasing in a statistical significant way, while
55
56 less significance is reached for the absolute yearly minimum temperature or the dates of
57
58 the first and last dates for the ensemble of the stations, may indicate some link to
59
60

1
2
3
4 changes in cloud cover. If the amount of clouds during night (or the number of cloudy
5
6 nights) increased, it might explain the rising of the minimum temperatures, whereas the
7
8 clear nights would cool as much as usual, thus allowing for similar yearly minimum
9
10 temperatures. The dates of first and last frost not significantly changed would then be
11
12 related to the fact that, whenever a clear and calm night occurs, similar intense cooling
13
14 as before can occur.
15
16
17

18
19
20 In table 4, the trends are computed for the humanly estimated cloud cover at 7, 13 and
21
22 18 local solar time plus the relative insolation extracted from the burned heliograph
23
24 bands. These measurements have a larger amount of uncertainty than temperatures, due
25
26 to the human factor in estimation of cloud cover and portion of band burned.
27
28
29
30
31
32
33
34
35
36
37

38 There is a statistically significant increase of clouds (low and total) at 7 local solar time,
39
40 a moment normally close to the time of the minimum temperature and a decrease of low
41
42 clouds at 13 and 18. This is combined with a small increase for the relative insolation.
43
44 This increase of late-night clouds would lead to higher minima for cloudy nights and
45
46 would be compatible with less significant trends for values usually found in clear and
47
48 calm nights, such as yearly minimum temperature. The increase of minimum
49
50 temperatures could thus be related, at least partially, to higher nocturnal thermal
51
52 stability, with more inversions -and therefore fog and low clouds-, while this stability
53
54 would also be compatible with a lower daily cloudiness and the corresponding increase
55
56 in relative sunshine hours.
57
58
59
60

1
2
3
4
5
6
7 The connection between these variables has been investigated by adjusting a multiple
8 regression model to the minimum daily temperatures, using cloudiness and relative
9 insolation as the independent variables. But first, all the variables have been detrended,
10 and their seasonal components removed by subtracting the annual cycle and the first
11 two harmonics of a Fourier analysis adjusted to the original data. In this way, the
12 variance of the minimum temperatures was reduced by a 75% (standard deviation of
13 6.3°C in the original data and 3.1°C in the deseasonalized one).
14
15
16
17
18
19
20
21
22
23
24
25

26 The residual series were then used to adjust the multiple regression model and, although
27 the explained variance of the minimum temperatures amounts only to 15%, only the low
28 cloudiness at 18 UTC was not significant at the 95% confidence level. The signs of the
29 significant contributions (Table 4) of the independent variables are in accordance with
30 the aforementioned interpretation of their trends.
31
32
33
34
35
36
37
38
39

40 **5. Conclusions**

41
42
43
44 For the ensemble of the studied stations located in the island of Majorca in the Western
45 Mediterranean, the analysis of the minimum temperatures for the last three decades
46 seems to imply that there is a statistically significant trend to increase, at about 0.6°C
47 per decade at a 95 % confidence level, more than the hemispheric estimations provided
48 by IPCC of about 0.3°C per decade . This goes together with a diminution of the number
49 of days of frost ($T < 0^{\circ}\text{C}$) or of cold ($T < 7^{\circ}\text{C}$). Instead, the absolute yearly minimum
50 temperature or the dates of the first and last frost days do not show a significant trend in
51
52
53
54
55
56
57
58
59
60

1
2
3
4 the ensemble. A possible link to cloud cover is considered, since this quantity has
5
6 significantly increased at 7 AM local solar time, that may partially explain the rise of
7
8 the minimum temperatures. In summary, warmer winters seem to be on place that may
9
10 induce relevant changes in the agricultural landscape of the region, but being aware that
11
12 frost events can take place as it has been usual in the last decades.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Peer Review Only

References

- Bartolini G, Morabito M, Crisci A, Grifoni D, Torrigiani T, Petralli, M, Marracchi G, Orlandini S. 2008. Recent trends in Tuscany (Italy) summer temperature and indices of extremes. *International Journal of Climatology* **28**: 1751-1760. DOI: 10.1002/joc.1673
- Cuxart J, Jimenez MA, Martinez D. 2007. Nocturnal Meso-Beta Basin and Katabatic Flows on a Midlatitude Island. *Monthly Weather Review* **135**: 918-932. DOI: 10.1175/MWR3329.1
- Easterling DR. 2002. Recent changes in frost days and the frost-free season in the United States. *Bulletin of the American Meteorological Society* **83**: 1327-1332
- Guijarro, JA. 2006. Homogenization of a dense thermo-pluviometric monthly database in the Balearic Islands using the free contributed R package CLIMATOL. *WMO Fifth Seminar for Homogenization and Quality Control in Climatological Databases*. Budapest, Hungary, 2006; WCDMP-No. 68, WMO-TD No. 1434, 2008.
- Jiménez MA, Mira A, Cuxart J, Luque A, Alonso S, Guijarro, JA. 2008. Verification of a clear-sky mesoscale simulation using satellite-derived surface temperatures. *Monthly Weather Review* **136**: 5148-5161. DOI: 10.1175/2008MWR2461.
- Kalma JD, Laughlin GP, Caprio JM, Hamer PJC. 1992. The bioclimatology of frost. Its occurrence, impact and protection. *Advances in Bioclimatology* **2**, pp 144, Springer-Verlag
- Kendall MG. 1975. Rank Correlation Methods, Charles Griffin, London
- Menzel A, Jakobi G, Ahas R, Scheifinger H, Estrella N. 2003. Variations of the climatological growing season (1951-2000) in Germany compared to other countries. *International Journal of Climatology* **23**: 793-812. DOI: 10.1002/joc.915
- R Development Core Team (2008). R: A language and environment for statistical

1
2
3
4 computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-
5
6
7 07-0, URL <http://www.R-project.org>.

8
9 Rayner, N.A., et al., 2006: Improved analyses of changes and uncertainties in sea
10
11 surface temperature measured *in situ* since the mid-nineteenth century: the HadSST2
12
13 dataset. *J. Clim.*, **19**, 446–469

14
15
16 Riera, A and Ripoll, AM, 2009: Informe econòmic i social de les Illes Balears, 2008,
17
18 Centre de Recerca Econòmica, UIB-Sa Nostra, Palma, Spain, ISBN 978-84-92863-00-6

19
20 Snyder R, de Melo Abreu J. 2005. Frost Protection: fundamentals, practice and
21
22 economics, vol 1. *Service Series 10, FAO, Roma*

23
24
25
26 Trenberth KE, Jones PD, Ambenje P, Bojariu R, Easterling D, Klein Tank A, Parker D,
27
28 Rahimzadeh F, Renwick JA, Rusticucci M, Soden B, Zhai P. 2007. Observations:
29
30 Surface and Atmospheric Climate Change. In: *Climate Change 2007: The Physical*
31
32 *Science Basis*. Contribution of Working Group I to the Fourth Assessment Report of the
33
34 Intergovernmental Panel on Climate Change [Solomon S, Qin D, Manning M, Chen Z,
35
36 Marquis M, Averyt KB, Tignor M, Miller HL(eds.)]. Cambridge University Press,
37
38 Cambridge, United Kingdom and New York, NY, USA.

39
40
41
42 Vose, R.S., D.R. Easterling, and B. Gleason, 2005: Maximum and minimum
43
44 temperature trends for the globe: An update through 2004. *Geophys. Res. Lett.*, **32**,
45
46 L23822, doi:10.1029/2004GL024379.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Tables

Table 1: Statistics of the daily and annual series of minimum temperatures for 8 selected Majorca stations. Average of the daily series is 0.60 C/decade. In bold regressions with confidence level higher than 95 %.

	Average Daily T min	Slope and 95% conf interval (C/decade)	Average annual T min	Slope and 95% conf. interval (C/ decade)
Palma Harbour (78-08) –Coast-	14.5 C	0.59 (0.48,0.71)	3.0 C	0.32 (-0.30, 0.94)
Sa Cabaneta (76-08) –Plateau-	13.1 C	0.62 (0.51,0.73)	1.2 C	0.45 (-0.18, 1.08)
Muro *(83-08) –Valley-	12.8 C	0.69 (0.52,0.85)	0.4 C	1.28 (0.62, 1.64)
Llucmajor (74-08) –Plateau-	12.6 C	0.34 (0.23,0.45)	1.2 C	0.03 (-0.45, 0.52)
Manacor* (84-08) –Valley-	11.8 C	0.70 (0.53,0.84)	-1.0 C	1.43 (-0.55, 2.31)
Ses Salines (72-08) – Valley-	10.5 C	0.31 (0.18,0.44)	-2.8 C	1.38 (0.61, 2.16)
Airport (72-08) –Valley--	10.3 C	0.64 (0.54,0.75)	-2.9 C	0.30 (-0.05, 0.65)
Orient (75-08) –Mountain-	8.4 C	0.99 (0.89,1.09)	-2.9 C	0.88 (0.37, 1.38)

Lluc (72-08)	8.0 C	0.55	-4.6 C	0.62
-Mountain-		(0.45,0.65)		(0.13, 1.11)

* after removal of parts of the series before the shifts in the mean: Muro (74-83),

Manacor (74-84)

Peer Review Only

Table 2: days with $T < 0$ (definition of day with frost). Day 1 is 1st January.

	Average Number of days	Slope 95% (per decade)	Average First day	Slope 95% (per decade)	Average Last day	Slope 95% (per decade)
Palma Harbour	0	N.A.	N.A.	N.A.	N.A.	N.A.
Sa Cabaneta	0.4	N.A.	N.A.	N.A.	N.A.	N.A.
Muro	0.8	N.A.	N.A.	N.A.	N.A.	N.A.
Llucmajor	0.1	N.A.	N.A.	N.A.	N.A.	N.A.
Manacor	0.8	N.A.	N.A.	N.A.	N.A.	N.A.
Ses Salines	13	(-8.8, -4.3)	-9 (23D)	(2.8, 17.2)	62 (3M)	(-28.9, -13.1)
Airport	13	(-5.4, -0.9)	-9 (23D)	(-1.5, 13.7)	70 (11M)	(-17.4, -3.8)
Orient	12	(-7.7, -1.4)	-2 (30D)	(-6.1, 20.5)	71 (12M)	(-21.6, -5.1)
Lluc	36	(-9.7, -0.2)	-29 (3D)	(-1.7, 11.6)	85 (26M)	(-12.0, 5.6)

Table 3: Number of days with $T < 7$ (definition of day with hours-of-cold). Day 1 is 1st January.

	Average number of days	Slope 95% (per decade)	Average first day	Slope 95% (per decade)	Average last day	Slope 95% (per decade)
Palma Harbour	28	(-13.5, -2.0)	-20 (12D)	(-9.8, 9.9)	78 (19M)	(-16.9, 2.9)
Sa Cabaneta	49	(-13.4, -0.5)	-33 (29N)	(-1.0, 11.0)	97 (7A)	(-15.6, -2.6)
Muro	72	(-21.4, -1.7)	-43 (19N)	(-2.8, 12.6)	104 (14A)	(-18.0, -3.8)
Llucmajor	62	(-5.6, 7.2)	-40 (22N)	(-5.2, 5.6)	100 (10A)	(-7.9, 0.4)
Manacor	91	(-23.8, -4.9)	-47 (15N)	(-9.4, 14.3)	115 (25A)	(-13.0, 2.3)
Ses Salines	110	(-29.7, -7.5)	-55 (7N)	(5.0, 17.2)	117 (27A)	(-35.4, -9.7)
Airport	127	(-16.8, -7.3)	-63 (29O)	(2.8, 10.5)	133 (12My)	(-13.5, -6.1)
Orient	145	(-33.7, -16.6)	-77 (15O)	(1.5, 11.1)	141 (12My)	(-14.9, -5.4)
Lluc	159	(-25.2, -3.8)	-80 (12O)	(-10.9, 1.6)	139 (18My)	(-13.6, 6.9)

Table 4: Averages (octas and percent) and trends (percent per decade) for cloudiness and relative insolation at Palma airport (1972-2008). Trends statistically significant at the 95% level are in bold. The third row displays the sign of the significant coefficients of a multiple regression between daily minimum temperature and these variables ("b" is the regression coefficient in $Var=a+bT_{min}$).

	Low clouds 7	Total clouds 7	Low clouds 13	Total clouds 13	Low clouds 18	Total clouds 18	Relative insolation
Average	2.0	3.7	2.3	3.9	1.8	3.6	60.8%
Trend	1.9%	2.2%	-2.4%	-0.3%	-1.7%	0.3%	1.3%
Sign of significant b	+	+	-	+	0	+	-

1
2
3
4 **Figure 1:** (a) Majorca in the Western Mediterranean; (b) Topographic map of the
5
6
7 Majorca island with the locations of the stations used.
8
9

10
11 **Figure 2.** *Top:* annual anomaly of the minimum temperature and absolute annual
12 minimum temperature, *second:* number of days below 0 C; *third:* number of days below
13 7 C; *fourth:* first day below 0°C and below 7 C; *bottom:* last day below 0 and below
14 7°C. Each figure has regression lines for the complete series.
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1: (a) Majorca in the Western Mediterranean
1561x995mm (72 x 72 DPI)





Figure 1: (b) Topographic map of the Majorca island with the locations of the stations used.
1594x1034mm (72 x 72 DPI)



207x268mm (150 x 150 DPI)